



Universidad  
de Murcia

Departamento  
Matemáticas

# Un teorema tipo Bishop-Phelps-Bollobás para álgebras uniformes

B. Cascales

<http://webs.um.es/beca>

I Encuentros AF Murcia-Almería  
10 de Mayo de 2013

Supported by



MINISTERIO  
DE ECONOMÍA  
Y COMPETITIVIDAD



Unión Europea  
Fondo Europeo de  
Desarrollo Regional

## Stay focused

- ③ **First part:** The need of the complex Urysohn type lemma.
  - Our result(s) in Functional Analysis needs the lemma.
  - A bit of history of the problems in FA above. People involved.

## Stay focused

- ③ **First part:** The need of the complex Urysohn type lemma.
  - Our result(s) in Functional Analysis needs the lemma.
  - A bit of history of the problems in FA above. People involved.
- ⑤ **Third part:** Applications (by others).

## Stay focused

- ③ **First part:** The need of the complex Urysohn type lemma.
  - Our result(s) in Functional Analysis needs the lemma.
  - A bit of history of the problems in FA above. People involved.
- ④ **Second part:** Complex Urysohn type lemma: a few pictures.
- ⑤ **Third part:** Applications (by others).



## Stay focused

- 2 The **topological ideas** behind the scenes.
- 3 **First part:** The need of the complex Urysohn type lemma.
  - Our result(s) in Functional Analysis needs the lemma.
  - A bit of history of the problems in FA above. People involved.
- 4 **Second part:** Complex Urysohn type lemma: a few pictures.
- 5 **Third part:** Applications (by others).

## Stay focused

- 1 **Co-authors.**
- 2 The **topological ideas** behind the scenes.
- 3 **First part:** The need of the complex Urysohn type lemma.
  - Our result(s) in Functional Analysis needs the lemma.
  - A bit of history of the problems in FA above. People involved.
- 4 **Second part:** Complex Urysohn type lemma: a few pictures.
- 5 **Third part:** Applications (by others).

# Co-authors

## Co-authors




R. M. Aron, B. Cascales and O. Kozhushkina,  
*The Bishop-Phelps-Bollobás theorem and Asplund operators*,  
Proc. Amer. Math. Soc. 139 (2011), no. 10, 3553–3560.



B. Cascales, A. J. Guirao and V. Kadets,  
*A Bishop-Phelps-Bollobás type theorem for uniform algebras*,  
Advances in Mathematics 240 (2013) 370–382

# Topology behind the scenes

## Topology behind the scenes.



$$\|\cdot\| - \text{diam}(U \cap S) \leq \varepsilon$$

## Asplund spaces: Namioka, Phelps and Stegall

Let  $X$  be a Banach space. Then the following conditions are equivalent:

- (i)  $X$  is an Asplund space, *i.e.*, whenever  $f$  is a convex continuous function defined on an open convex subset  $U$  of  $X$ , the set of all points of  $U$  where  $f$  is Fréchet differentiable is a dense  $G_\delta$ -subset of  $U$ .
- (ii) every  $w^*$ -compact subset of  $(X^*, w^*)$  is fragmented by the norm;
- (iii) each separable subspace of  $X$  has separable dual;
- (iv)  $X^*$  has the Radon-Nikodým property.



Our result(s) in functional analysis: the need of the lemma

Complex Urysohn type lemma: a few pictures

Applications

Presentation

Bishop-Phelps-Bollobás property

A bit of history of the problems in FA. People involved.

Urysohn's lemma plays its part

An idea of the proof for  $\mathfrak{A} = A(\mathbb{D})$

# Our result(s) in functional analysis: the need of the lemma



## Our result(s)

Theorem (A. J. Guirao, V. Kadets and B. C. 2012)

Let  $\mathfrak{A} \subset C(K)$  be a uniform algebra and  $T: X \rightarrow \mathfrak{A}$  be an Asplund operator with  $\|T\| = 1$ . Suppose that  $0 < \varepsilon < \sqrt{2}$  and  $x_0 \in S_X$  are such that  $\|Tx_0\| > 1 - \frac{\varepsilon^2}{2}$ . Then there exist  $u_0 \in S_X$  and an Asplund operator  $\tilde{T} \in S_{L(X, \mathfrak{A})}$  satisfying that

$$\|\tilde{T}u_0\| = 1, \|x_0 - u_0\| \leq \varepsilon \quad \text{and} \quad \|T - \tilde{T}\| < 2\varepsilon.$$

Theorem (R. M. Aron, O. Kozhushkina and B. C. 2011)

Let  $T: X \rightarrow C(K)$  be an Asplund operator with  $\|T\| = 1$ . Suppose that  $0 < \varepsilon < \sqrt{2}$  and  $x_0 \in S_X$  are such that  $\|Tx_0\| > 1 - \frac{\varepsilon^2}{2}$ . Then there exist  $u_0 \in S_X$  and an Asplund operator  $\tilde{T} \in S_{L(X, C(K))}$  satisfying that

$$\|\tilde{T}u_0\| = 1, \|x_0 - u_0\| \leq \varepsilon \quad \text{and} \quad \|T - \tilde{T}\| < 2\varepsilon.$$

## Recall that...

### Uniform algebra

- 1 Closed vector subspace  $A \subset C(K)$ , with the properties:
  - the products of functions in  $A$  remains in  $A$ ;
  - either  $1 \in A$  or  $A$  distinguishes the points of  $K$ .
  - $A$  separates the points of  $K$ .

## Recall that...

### Uniform algebra

- 1 Closed vector subspace  $A \subset C(K)$ , with the properties:
  - the products of functions in  $A$  remains in  $A$ ;
  - either  $1 \in A$  or  $A$  distinguishes the points of  $K$ .
  - $A$  separates the points of  $K$ .
- 2 In the real case, if  $A$  uniform algebra  $\Rightarrow A = C(K)$ .

## Recall that...


### Uniform algebra

- 1 Closed vector subspace  $A \subset C(K)$ , with the properties:
  - the products of functions in  $A$  remains in  $A$ ;
  - either  $1 \in A$  or  $A$  distinguishes the points of  $K$ .
  - $A$  separates the points of  $K$ .
- 2 In the real case, if  $A$  uniform algebra  $\Rightarrow A = C(K)$ .
- 3 The complex case is **different**: the disk algebra  $A(\mathbb{D})$ .

## Asplund operators

Stegall, 1975

An **operator**  $T \in L(X, Y)$  is **Asplund**, if it factors through an Asplund space:




$Z$  is Asplund;  $T_1 \in L(X, Z)$  and  $T_2 \in L(Z, Y)$ .

$T$  Asplund operator  $\Leftrightarrow T^*(B_{Y^*})$  is fragmented by the norm of  $X^*$ .

# Asplund operators

Stegall, 1975

An **operator**  $T \in L(X, Y)$  is **Asplund**, if it factors through an Asplund space:



$Z$  is Asplund;  $T_1 \in L(X, Z)$  and  $T_2 \in L(Z, Y)$ .

$T$  Asplund operator  $\Leftrightarrow T^*(B_{Y^*})$  is fragmented by the norm of  $X^*$ .

Theorem (A. J. Guirao, V. Kadets and B. C. 2012)

Let  $\mathfrak{A} \subset C(K)$  be a uniform algebra and  $T: X \rightarrow \mathfrak{A}$  be an Asplund operator with  $\|T\| = 1$ . Suppose that  $0 < \varepsilon < \sqrt{2}$  and  $x_0 \in S_X$  are such that

$\|Tx_0\| > 1 - \frac{\varepsilon^2}{2}$ . Then there exist  $u_0 \in S_X$  and an Asplund operator  $\tilde{T} \in S_{L(X, \mathfrak{A})}$  satisfying that

$$\|\tilde{T}u_0\| = 1, \|x_0 - u_0\| \leq \varepsilon$$

and

$$\|T - \tilde{T}\| < 2\varepsilon.$$

### Corollary

Let  $T \in L(X, \mathfrak{A})$  **weakly compact** with  $\|T\| = 1$ ,  $\frac{1}{2} > \varepsilon > 0$ , and  $x_0 \in S_X$  be such that

$$\|T(x_0)\| > 1 - \frac{\varepsilon^2}{4}.$$

Then there are  $u_0 \in S_X$  and  $S \in L(X, \mathfrak{A})$  **weakly compact** with  $\|S\| = 1$  satisfying

$$\|S(u_0)\| = 1, \|x_0 - u_0\| < \varepsilon \text{ and } \|T - S\| \leq 2\varepsilon.$$

### Corollary

$(X, \mathfrak{A})$  has the BPBP for any Asplund space  $X$  and any locally compact Hausdorff topological space  $L$  ( $X = c_0(\Gamma)$ , **for instance**).

### Corollary

$(X, C_0(L))$  has the BPBP for any  $X$  and any scattered locally compact Hausdorff topological space  $L$ .

Theorem (A. J. Guirao, V. Kadets and B. C. 2012)

Let  $\mathfrak{A} \subset C(K)$  be a uniform algebra and  $T: X \rightarrow \mathfrak{A}$  be an Asplund operator with  $\|T\| = 1$ . Suppose that  $0 < \varepsilon < \frac{1}{\sqrt{2}}$  and  $x_0 \in S_X$  are such that  $\|Tx_0\| > 1 - \frac{\varepsilon^2}{2}$ . Then there exist  $u_0 \in S_X$  and an Asplund operator  $\tilde{T} \in S_{L(X, \mathfrak{A})}$  satisfying that

$$\|\tilde{T}u_0\| = 1, \|x_0 - u_0\| \leq \varepsilon$$

and

$$\|T - \tilde{T}\| < 2\varepsilon.$$

## Corollary

Let  $T \in L(X, A(\mathbb{D}))$  weakly compact with  $\|T\| = 1$ ,  $\frac{1}{2} > \varepsilon > 0$ , and  $x_0 \in S_X$  be such that

$$\|Tx_0\| > 1 - \frac{\varepsilon^2}{4}.$$

Then there are  $u_0 \in S_X$  and  $S \in L(X, A(\mathbb{D}))$  weakly compact with  $\|S\| = 1$  satisfying

$$\|S(u_0)\| = 1, \|x_0 - u_0\| < \varepsilon \text{ and } \|T - S\| \leq 2\varepsilon.$$

## Remark

The theorem applies in particular to the ideals of finite rank operators  $\mathcal{F}$ , compact operators  $\mathcal{K}$ ,  $p$ -summing operators  $\Pi_p$  and of course to the weakly compact operators  $\mathcal{W}$  themselves. To the best of our knowledge even in the case  $\mathcal{W}(X, \mathfrak{A})$  the Bishop-Phelps property that follows is a brand new result.



## Bishop-Phelps theorem

### Theorem (Bishop-Phelps, 1961)

If  $X$  is a Banach, then  $\overline{\mathbf{N}AX^*} = X^*$ .

### A PROOF THAT EVERY BANACH SPACE IS SUBREFLEXIVE

BY ERRETT BISHOP AND R. R. PHELPS

Communicated by Mahlon M. Day, August 19, 1960

A real or complex normed space is *subreflexive* if those functionals which attain their supremum on the unit sphere  $S$  of  $E$  are norm-dense in  $E^*$ , i.e., if for each  $f$  in  $E^*$  and each  $\epsilon > 0$  there exist  $g$  in  $E^*$  and  $x$  in  $S$  such that  $|g(x)| = \|g\|$  and  $\|f - g\| < \epsilon$ . There exist incomplete normed spaces which are not subreflexive [1]<sup>1</sup> as well as incomplete spaces which *are* subreflexive (e.g., a dense subspace of a Hilbert space). It is evident that every reflexive Banach space is sub-

# The Bishop-Phelps property for operators

## Definition

An operator  $T : X \rightarrow Y$  is **norm attaining** if there exists  $x_0 \in X$ ,  $\|x_0\| = 1$ , such that  $\|T(x_0)\| = \|T\|$ .

## Definition (Lindenstrauss)

$(X, Y)$  has the Bishop-Phelps Property (BPP) if every operator  $T : X \rightarrow Y$  can be uniformly approximated by **norm attaining** operators.

- 1  $(X, \mathbb{K})$  has BPP for every  $X$   
**Bishop-Phelps** (1961);
- 2  $\overline{\{T \in L(X; Y) : T^{**} \in NA(X^{**}; Y^{**})\}} = L(X; Y)$  for every pair of Banach spaces  $X$  and  $Y$ , **Lindenstrauss** (1963);
- 3  $X$  with RNP, then  $(X, Y)$  has BPP for every  $Y$ , **Bourgain** (1977);
- 4 there are spaces  $X, Y$  and  $Z$  such that  $(X, C([0, 1]))$ ,  $(Y, \ell^p)$  ( $1 < p < \infty$ ) and  $(Z, L^1([0, 1]))$  fail BPP, **Schachermayer** (1983), **Gowers** (1990) and **Acosta** (1999);
- 5  $(C(K), C(S))$  has BPP for all compact spaces  $K, S$ , **Johnson and Wolfe**, (1979).
- 6  $(L^1([0, 1]), L^\infty([0, 1]))$  has BPP,  
**Finet-Payá** (1998).

## Bollobás observation, 1970

### AN EXTENSION TO THE THEOREM OF BISHOP AND PHELPS

BÉLA BOLLOBÁS

Bishop and Phelps proved in [1] that every real or complex Banach space is *subreflexive*, that is the functionals (real or complex) which attain their supremum on the unit sphere of the space are dense in the dual space. We shall sharpen this result and then apply it to a problem about the numerical range of an operator.

Denote by  $S$  and  $S'$  the unit spheres of  $X$  and  $X^*$  respectively.

**THEOREM 1.** *Suppose  $X$  is a Banach space. If  $S$  and  $S'$  are such that for every  $y \in S$  and  $g \in S'$  such*

Corollary... the way it is oftentimes presented

Given  $\frac{1}{2} > \varepsilon > 0$ , if  $x_0 \in S_X$  and  $x^* \in S_{X^*}$  are such that

$$|x^*(x_0)| > 1 - \frac{\varepsilon^2}{4},$$

then there are  $u_0 \in S_X$  and  $y^* \in S_{X^*}$  such that

$$|y^*(u_0)| = 1, \|x_0 - u_0\| < \varepsilon \text{ and } \|x^* - y^*\| < \varepsilon.$$

# Bishop-Phelps-Bollobás Property for operators

**Definition:** Acosta, Aron, García and Maestre, 2008

$(X, Y)$  is said to have the Bishop-Phelps-Bollobás property (BPBP) if for any  $\varepsilon > 0$  there are  $\eta(\varepsilon) > 0$  such that for all  $T \in S_{L(X, Y)}$ , if  $x_0 \in S_X$  is such that

$$\|T(x_0)\| > 1 - \eta(\varepsilon),$$

then there are  $u_0 \in S_X$ ,  $S \in S_{L(X, Y)}$  with

$$\|S(u_0)\| = 1$$

and

$$\|x_0 - u_0\| < \varepsilon \text{ and } \|T - S\| < \varepsilon.$$

- 1  $Y$  has *certain* almost-biorthogonal system  $(X, Y)$  has BPBP any  $X$ ;
- 2  $(\ell^1, Y)$  BPBP is characterized through a condition called AHSP: it holds for  $Y$  finite dimensional, uniformly convex,  $Y = L^1(\mu)$  for a  $\sigma$ -finite measure or  $Y = C(K)$ ;
- 3 there is pair  $(\ell^1, X)$  failing BPBP, but having BPBP;
- 4  $(\ell_n^\infty, Y)$  has BPBP  $Y$  uniformly convex **no hope for  $c_0$** :  
 $\eta(\varepsilon) = \eta(n, \varepsilon) \rightarrow 1$  with  $n \rightarrow \infty$ .

## Bishop-Phelps-Bollobás Property for operators

**Definition:** Acosta, Aron, García and Maestre, 2008

$(X, Y)$  is said to have the Bishop-Phelps-Bollobás property (BPBP) if for any  $\varepsilon > 0$  there are  $\eta(\varepsilon) > 0$  such that for all  $T \in S_{L(X, Y)}$ , if  $x_0 \in S_X$  is such that

$$\|T(x_0)\| > 1 - \eta(\varepsilon),$$

then there are  $u_0 \in S_X$ ,  $S \in S_{L(X, Y)}$  with

$$\|S(u_0)\| = 1$$

and

$$\|x_0 - u_0\| < \varepsilon \text{ and } \|T - S\| < \varepsilon.$$

- 1  $Y$  has *certain* almost-biorthogonal system  $(X, Y)$  has BPBP any  $X$ ;
- 2  $(\ell^1, Y)$  BPBP is characterized through a condition called AHSP: it holds for  $Y$  finite dimensional, uniformly convex,  $Y = L^1(\mu)$  for a  $\sigma$ -finite measure or  $Y = C(K)$ ;
- 3 there is pair  $(\ell^1, X)$  failing BPBP, but having BPBP;
- 4  $(\ell_n^\infty, Y)$  has BPBP  $Y$  uniformly convex **no hope for  $c_0$** :  
 $\eta(\varepsilon) = \eta(n, \varepsilon) \rightarrow 1$  with  $n \rightarrow \infty$ .

### PROBLEM?

No  $Y$  infinite dimensional was known s.t.  $(c_0, Y)$  has BPBP.

# An idea of the proof for $\mathfrak{A} = C(K)$

## Theorem (R. M. Aron, O. Kozhushkina and B. C. 2011)

Let  $T: X \rightarrow C(K)$  be an Asplund operator with  $\|T\| = 1$ . Suppose that  $0 < \varepsilon < \sqrt{2}$  and  $x_0 \in S_X$  are such that  $\|Tx_0\| > 1 - \frac{\varepsilon^2}{2}$ . Then there exist  $u_0 \in S_X$  and an Asplund operator  $\tilde{T} \in S_{L(X, C(K))}$  satisfying that

$$\|\tilde{T}u_0\| = 1, \|x_0 - u_0\| \leq \varepsilon \quad \text{and} \quad \|T - \tilde{T}\| < 2\varepsilon.$$

- 1 **Topological tools** provide a **suitable** open set  $U \subset K$ ,  $y^* \in S_{X^*}$  and  $\rho < 2\varepsilon$  with

$$1 = |y^*(u_0)| = \|u_0\| \quad \text{and} \quad \|x_0 - u_0\| < \varepsilon \quad \& \quad \|T^*(\delta_t) - y^*\| < \rho \quad \forall t \in U$$

- 2 **Urysohn's lemma** that applied to an arbitrary  $t_0 \in U$  produces a function  $f \in C(K)$  satisfying



$$f(t_0) = \|f\|_\infty = 1, f(K) \subset [0, 1] \quad \text{and} \quad \text{supp}(f) \subset U.$$

- 3  $\tilde{T}$  is explicitly defined by

$$\tilde{T}(x)(t) = f(t) \cdot y^*(x) + (1 - f(t)) \cdot T(x)(t), \quad x \in X, t \in K,$$

- 4 The **suitability** of  $U$  is used to prove that  $\|T - \tilde{T}\| < 2\varepsilon$ .

# A few words about Urysohn's lemma



## Normal spaces - Exercise for students

For a topological space  $T$  the following statements are equivalent:

- 1  $T$  is normal.
- 2 Urysohn's lemma holds for  $T$ .
- 3 Tietze's extension theorem holds for  $T$ .
- 4 Katetov's "sandwich" theorem holds for  $T$ .
- 5 For every function  $f \in \mathbb{R}^T$  the distance to  $C_b(T)$  is given by

$$d(f, C_b(T)) = \frac{1}{2} \text{osc}(f).$$

# An idea of the proof for $\mathfrak{A} = A(\mathbb{D})$

## Theorem (A. J. Guirao, V. Kadets and B. C. 2012)

Let  $T: X \rightarrow A(\mathbb{D})$  be an Asplund operator with  $\|T\| = 1$ . Suppose that  $0 < \varepsilon < \sqrt{2}$  and  $x_0 \in S_X$  are such that  $\|Tx_0\| > 1 - \frac{\varepsilon^2}{2}$ . Then there exist  $u_0 \in S_X$  and an Asplund operator  $\tilde{T} \in S_{L(X, A(\mathbb{D}))}$  satisfying that

$$\|\tilde{T}u_0\| = 1, \|x_0 - u_0\| \leq \varepsilon \quad \text{and} \quad \|T - \tilde{T}\| < 2\varepsilon.$$


- Topological tools** gives an open set  $U \subset \mathbb{D}$   $y^* \in S_{X^*}$  &  $\rho < 2\varepsilon$  with  $1 = |y^*(u_0)| = \|u_0\|$  and  $\|x_0 - u_0\| < \varepsilon$  &  $\|T^*(\delta_t) - y^*\| < \rho \forall t \in U$ .
- Urysohn's lemma that applied to an arbitrary  $t_0 \in U$  produces a function  $f \in A(\mathbb{D})$  satisfying

$$f(t_0) = \|f\|_\infty = 1, f(\mathbb{D}) \subset [0, 1] \quad \text{and} \quad \text{supp}(f) \subset U.$$

explicitly defined by

$$\tilde{T}(x)(t) = f(t) \cdot y^*(x) + (1 - f(t)) \cdot T(x)(t), \quad x \in X, t \in \mathbb{D},$$

continuity of  $U$  is used to prove that  $\|T - \tilde{T}\| < 2\varepsilon$ .





## An idea of the proof for $\mathfrak{A} = A(\mathbb{D})$

### Theorem (A. J. Guirao, V. Kadets and B. C. 2012)

Let  $T: X \rightarrow A(\mathbb{D})$  be an Asplund operator with  $\|T\| = 1$ . Suppose that  $0 < \varepsilon < \sqrt{2}$  and  $x_0 \in S_X$  are such that  $\|Tx_0\| > 1 - \frac{\varepsilon^2}{2}$ . Then there exist  $u_0 \in S_X$  and an Asplund operator  $\tilde{T} \in S_{L(X, A(\mathbb{D}))}$  satisfying that

$$\|\tilde{T}u_0\| = 1, \|x_0 - u_0\| \leq \varepsilon \quad \text{and} \quad \|T - \tilde{T}\| < 2\varepsilon.$$

1 **Topological tools** gives an open set,  $U \cap \mathbb{T} \neq \emptyset$ ,  $y^* \in S_{X^*}$  &  $\rho < 2\varepsilon$  with  
 $1 = |y^*(u_0)| = \|u_0\|$  and  $\|x_0 - u_0\| < \varepsilon$  &  $\|T^*(\delta_t) - y^*\| < \rho \forall t \in U$ .

2 Urysohn's lemma that applied to an arbitrary  $t_0 \in U \cap \mathbb{T}$  produces a function  $f \in A(\mathbb{D})$  satisfying

$$f(t_0) = \|f\|_\infty = 1, f(\overline{\mathbb{D}}) \subset R_{\varepsilon'} \quad \text{and} \quad f \text{ small in } \overline{\mathbb{D}} \setminus U.$$

3  $\tilde{T}$  is explicitly defined by

$$\tilde{T}(x)(t) = f(t) \cdot y^*(x) + (1 - \varepsilon')(1 - f(t)) \cdot T(x)(t)$$

4 The suitability of  $U$  is used to prove that  $\|T - \tilde{T}\| < 2\varepsilon$ .

# An idea of the proof for $\mathfrak{A} = A(\mathbb{D})$

## Theorem (A. J. Guirao, V. Kadets and B. C. 2012)

Let  $T: X \rightarrow A(\mathbb{D})$  be an Asplund operator with  $\|T\| = 1$ . Suppose that  $0 < \varepsilon < \sqrt{2}$  and  $x_0 \in S_X$  are such that  $\|Tx_0\| > 1 - \frac{\varepsilon^2}{2}$ . Then there exist  $u_0 \in S_X$  and an Asplund operator  $\tilde{T} \in S_{L(X, A(\mathbb{D}))}$  satisfying that

$$\|\tilde{T}u_0\| = 1, \|x_0 - u_0\| \leq \varepsilon \quad \text{and} \quad \|T - \tilde{T}\| < 2\varepsilon.$$

- Topological tools** gives an open set,  $U \cap \mathbb{T} \neq \emptyset$ ,  $y^* \in S_{X^*}$  &  $\rho < 2\varepsilon$  with  $1 = |y^*(u_0)| = \|u_0\|$  and  $\|x_0 - u_0\| < \varepsilon$  &  $\|T^*(\delta_t) - y^*\| < \rho \forall t \in U$ .
- Urysohn's lemma that applied to an arbitrary  $t_0 \in U \cap \mathbb{T}$  produces a function  $f \in A(\mathbb{D})$  satisfying


$$f(t_0) = \|f\|_\infty = 1, f(\overline{\mathbb{D}}) \subset R_{\varepsilon'}$$

and  $f$  small in  $\overline{\mathbb{D}} \setminus U$ .

explicitly defined by

$$\tilde{T}(x)(t) = f(t) \cdot y^*(x) + (1 - \varepsilon')(1 - f(t)) \cdot T(x)(t)$$

suitability of  $U$  is used to prove that  $\|T - \tilde{T}\| < 2\varepsilon$ .





# Complex Urysohn type lemma: a few pictures

Urysohn type lemma for  $A(\mathbb{D})$ 

$U \subset^{\text{open}} \mathbb{D}$ ,  $1 \in U$ ,  $\exists f(1) = \|f\|_{\infty} = 1$ ,  $f(\mathbb{D}) \subset R_{\varepsilon'}$  and  $f$  small in  $\mathbb{D} \setminus U$ ,

$$f_n(z) := \left(\frac{z+1}{2}\right)^n$$







## Urysohn type lemma for $A(\mathbb{D})$

$U \subset^{\text{open}} \overline{\mathbb{D}}$ ,  $1 \in U$ ,  $\exists f(1) = \|f\|_{\infty} = 1$ ,  $f(\overline{\mathbb{D}}) \subset R_{\varepsilon'}$  and  $f$  small in  $\overline{\mathbb{D}} \setminus U$ ,

$$f_n(z) := \left(\frac{z+1}{2}\right)^n$$

$$\exists n_1, n_2, \dots, n_k$$

$$f := \frac{f_{n_1} + f_{n_2} + \dots + f_{n_k}}{k}$$






## Uryshon type lemma for $A(\mathbb{D})$

$U \subset^{open} \overline{\mathbb{D}}, 1 \in U, \exists f(1) = \|f\|_\infty = 1, f(\overline{\mathbb{D}}) \subset R_{\varepsilon'}$  and  $f$  small in  $\overline{\mathbb{D}} \setminus U$ ,

$$z \rightarrow \frac{z + \delta}{1 + \delta}$$




## Urysohn type lemma for $A(\mathbb{D})$



$U \subset \overset{\text{open}}{\mathbb{D}}, 1 \in U, \exists f(1) = \|f\|_\infty = 1, f(\overline{\mathbb{D}}) \subset R_{\varepsilon'}$  and  $f$  small in  $\overline{\mathbb{D}} \setminus U$ ,


$$z \rightarrow 1 - \sqrt{1 - z}$$



Urysohn type lemma for  $A(\mathbb{D})$ 

$U \subset \overline{\mathbb{D}}$ ,  $1 \in U$ ,  $\exists f(1) = \|f\|_\infty = 1$ ,  $f(\overline{\mathbb{D}}) \subset R_{\varepsilon'}$  and  $f$  small in  $\overline{\mathbb{D}} \setminus U$ ,

## Riemann conformal mapping



Our Urysohn type lemma is suited for calculations with a computer.





## Why can we use the computer? Because of the provided proofs.

$$f_n(z) := \left(\frac{z+1}{2}\right)^n$$

$$\exists n_1, n_2, \dots, n_k$$

$$f := \frac{f_{n_1} + f_{n_2} + \dots + f_{n_k}}{k}$$



*Proof.* Let us fix  $0 < \eta < \min\{\delta/6, 1/2\}$  and  $n \in \mathbb{N}$ ,  $n > \frac{2}{\eta}$ . Let us define  $U_1 := U$ . We shall construct inductively a collection of points  $\{t_j\}_{j=1}^n$ , a decreasing finite sequence  $\{U_j\}_{j=1}^{n+1}$  of open subsets of  $U$ ,  $t_j \in U_{j+1} \cap \Gamma_0$ , and functions  $\{f_j\}_{j=1}^n \subset A$ , satisfying for any  $j \in \{1, \dots, n\}$  the following conditions:

- (i)  $f_j(t_j) = 1$ .
- (ii)  $|f_j(t)| < \frac{\eta}{2}$  for  $t \in K \setminus U_j$ .
- (iii)  $|f_j(t) - 1| < \frac{\eta}{2}$  for  $t \in U_{j+1}$ .

Indeed, Lemma 2.1 allows us to find a norm one function  $f_1 \in A$  and a  $t_1 \in U_1 \cap \Gamma_0$  such that  $f_1(t_1) = 1$  and  $|f_1(t)| < \frac{\eta}{2}$  for  $t \in K \setminus U_1$ . **THE PROOF GOES ON**

...

## 1+4+1 applications

- (1) The one already presented for the BPBp by Guirao-Kadets-Cascales;
- In the paper

SOME GEOMETRIC PROPERTIES ON DISK ALGEBRAS

YUN SUNG CHOI, DOMINGO GARCÍA, SUN KWANG KIM AND MANUEL MAESTRE

- (2) Daugavet prop.;
- (3) Lusheness;
- (4) Numerical radius;
- (5) AHSP.
- (6) in the Ph. dissertation by O. Kozhushkina for BPBp for  $\mathfrak{A}(K, Y)$

**Thank you**