

Domination by second countable spaces and Lindelöf Σ -property

B. CASCALES^{1,2}, J. ORIHUELA^{1,2} AND V.V. TKACHUK^{3,4}

Abstract. Given a space M , a family of sets \mathcal{A} of a space X is *ordered by M* if $\mathcal{A}=\{A_K:K \text{ is a compact subset of } M\}$ and $K \subset L$ implies $A_K \subset A_L$. We study the class \mathcal{M} of spaces which have compact covers ordered by a second countable space. We prove that a space $C_p(X)$ belongs to \mathcal{M} if and only if it is a Lindelöf Σ -space. Under $\text{MA}(\omega_1)$, if X is compact and $(X \times X) \setminus \Delta$ has a compact cover ordered by a Polish space then X is metrizable; here $\Delta = \{(x,x):x \in X\}$ is the diagonal of the space X . Besides, if X is a compact space of countable tightness and $X^2 \setminus \Delta$ belongs to \mathcal{M} then X is metrizable in ZFC.

We also consider the class \mathcal{M}^* of spaces X which have a compact cover \mathcal{F} ordered by a second countable space with the additional property that, for every compact set $P \subset X$ there exists $F \in \mathcal{F}$ with $P \subset F$. It is a ZFC result that if X is a compact space and $(X \times X) \setminus \Delta$ belongs to \mathcal{M}^* then X is metrizable. We also establish that, under CH, if X is compact and $C_p(X)$ belongs to \mathcal{M}^* then X is countable.

Keywords: (strong) domination by irrationals, (strong) domination by a second countable space, diagonal, metrization, orderings by irrationals, orderings by a second countable space, compact cover, function spaces, cosmic spaces, \aleph_0 -spaces, Lindelöf Σ -space, compact space, metrizable space

2000 Mathematics Subject Classification: 54B10, 54C05, 54D30

0. Introduction.

Given a space X we denote by $\mathcal{K}(X)$ the family of all compact subsets of X . One of about a dozen equivalent definitions says that X is a *Lindelöf Σ -space (or has the Lindelöf Σ -property)* if there exists a second countable space M and a compact-valued upper semicontinuous map $\varphi : M \rightarrow X$ such that $\bigcup \{\varphi(x) : x \in M\} = X$ (see, e.g., [RJ, Section 5.1]). It is worth mentioning that in Functional Analysis, the same concept is usually referred to as a *countably K -determined space*.

Suppose that X is a Lindelöf Σ -space and hence we can find a compact-valued upper semicontinuous surjective map $\varphi : M \rightarrow X$ for some second countable space M . If we let $F_K = \bigcup \{\varphi(x) : x \in K\}$ for any compact set $K \subset M$ then the family $\mathcal{F} = \{F_K : K \in \mathcal{K}(M)\}$ consists of compact subsets of X , covers X and $K \subset L$ implies $F_K \subset F_L$. We will say that \mathcal{F} is an M -ordered compact cover of X .

The class \mathcal{M} of spaces with an M -ordered compact cover for some second countable space M , was introduced by Cascales and Orihuela in [CO2]. They proved, among other things, that a Dieudonné complete space is Lindelöf Σ if and

¹ Research supported by FEDER and MEC, Project MTM2008-05396

² Research supported by Fundación Séneca de la CARM, Project 08848/PI/08

³ Research supported by Consejo Nacional de Ciencia y Tecnología de México, Grant U48602-F

⁴ Research supported by Programa Integral de Fortalecimiento Institucional (PIFI), Grant 34536-55

only if it belongs to \mathcal{M} . We proved in the previous paragraph that any Lindelöf Σ -space belongs to \mathcal{M} ; however, $X \in \mathcal{M}$ does not even imply that X is Lindelöf (see [CO2] and [Tk]) so \mathcal{M} is a new class which seems to be interesting in itself.

Let \mathbb{P} be the set of the irrationals which we will identify with ω^ω ; a family \mathcal{A} of subsets of a space X is \mathbb{P} -directed if $\mathcal{A} = \{A_p : p \in \mathbb{P}\}$ and $p \leq q$ implies $A_p \subset A_q$. The spaces which have \mathbb{P} -directed compact covers were extensively studied in Functional Analysis (see [Ca], [CO1], [FS1], [FS2], [KS] and [Ta]). Talagrand proved in [Ta] that if X is compact then $C_p(X)$ has a \mathbb{P} -directed compact cover if and only if $C_p(X)$ is K -analytic. Cascales [Ca] extended Talagrand's results by proving that, for angelic spaces, to have a \mathbb{P} -directed compact cover is equivalent to K -analyticity. Tkachuk [Tk] studied systematically the topology of the spaces which have a \mathbb{P} -directed compact cover (calling the respective spaces \mathbb{P} -dominated); it was proved in [Tk] that compactness can be omitted in the mentioned Talagrand result, i.e., for any Tychonoff X , the space $C_p(X)$ is K -analytic if and only if it is \mathbb{P} -dominated.

Following the terminology of [Tk] we say that a space X is M -dominated (or dominated by space M) if X has an M -ordered compact cover, i.e., there exists a family $\mathcal{F} = \{F_K : K \in \mathcal{K}(M)\} \subset \mathcal{K}(X)$ such that $\bigcup \mathcal{F} = X$ and $K \subset L$ implies $F_K \subset F_L$ for any $K, L \in \mathcal{K}(M)$. In this paper we study the general topological and categorical properties of the class \mathcal{M} of spaces dominated by a second countable space.

We prove, in particular, that for any Tychonoff X , the space $C_p(X)$ has the Lindelöf Σ -property, if and only if it is dominated by a second countable space. We also show that, if X is a compact space of countable tightness and $(X \times X) \setminus \Delta$ belongs to the class \mathcal{M} then X is metrizable. Here $\Delta = \{(x, x) : x \in X\}$ is the diagonal of the space X . It turns out that, under $\text{MA}(\omega_1)$, if X is compact and $(X \times X) \setminus \Delta$ is dominated by a Polish space then X is metrizable. As in [Tk], we introduce the notion of a strong M -domination to prove that if X is compact and the space $(X \times X) \setminus \Delta$ is strongly dominated by a second countable space then X is metrizable. Besides, under the continuum Hypothesis (CH), if X is compact and $C_p(X)$ is strongly dominated by a second countable space then X is countable. Hopefully, our study of M -dominated spaces will find applications in Functional Analysis the same as \mathbb{P} -domination already did.

1. Notation and terminology.

All spaces under consideration are assumed to be Tychonoff. If X is a space then $\tau(X)$ is its topology and $\tau^*(X) = \tau(X) \setminus \{\emptyset\}$. If X is a space and $A \subset X$ then $\tau(A, X) = \{U \in \tau(X) : A \subset U\}$; we will write $\tau(x, X)$ instead of $\tau(\{x\}, X)$. Given a space Z the family $\mathcal{K}(Z)$ consists of all compact subsets of Z ; we use the symbol \mathbb{P} to denote the set of the irrational numbers which we identify with ω^ω . Given $p, q \in \mathbb{P}$ we write $p \leq q$ if $p(n) \leq q(n)$ for any $n \in \omega$; we use the notation $p \leq^* q$ (or $p =^* q$) if there exists $m \in \omega$ such that $p(n) \leq q(n)$ (or $p(n) = q(n)$ respectively)

for all $n \geq m$. The symbol \mathbb{Q} stands for the set of the rational numbers with the topology induced from the real line \mathbb{R} and $\mathbb{N} = \omega \setminus \{0\}$.

A family of sets \mathcal{A} is \mathbb{P} -directed if $\mathcal{A} = \{A_p : p \in \mathbb{P}\}$ and $p \leq q$ implies $A_p \subset A_q$. A family \mathcal{B} is M -ordered for some space M if $\mathcal{B} = \{B_K : K \in \mathcal{K}(M)\}$ while $K \subset L$ implies $B_K \subset B_L$. A space X is \mathbb{P} -dominated if it has a \mathbb{P} -ordered compact cover; in general, the space X is dominated by a space M if it has an M -ordered compact cover. Say that X is *strongly M -dominated* if it has an M -ordered compact cover \mathcal{C} such that for every compact subset $K \subset X$ there exists $C \in \mathcal{C}$ with $K \subset C$.

If X is a space and \mathcal{C} is a cover of X then a family \mathcal{F} is called a *network modulo \mathcal{C}* if for any $C \in \mathcal{C}$ and $U \in \tau(C, X)$ there is $F \in \mathcal{F}$ with $C \subset F \subset U$. A family \mathcal{N} of subsets of a space X is a network in X if it is a network modulo the cover $\{\{x\} : x \in X\}$. The network weight $nw(X)$ of a space X is the minimal cardinality of a network in X . A space X is *cosmic* if $nw(X) = \omega$.

A cover \mathcal{C} of X is compact if all elements of \mathcal{C} are compact. A space X is *Lindelöf Σ* if it has a countable network modulo a compact cover of X . Say that X is an \aleph_0 -space if it has a countable network modulo $\mathcal{K}(X)$. The space X is *hemicompact* if there exists a countable family \mathcal{F} of compact subsets of X such that every $K \in \mathcal{K}(X)$ is contained in an element of \mathcal{F} .

If X is a space then $\Delta = \{(x, x) : x \in X\}$ is its diagonal. The space X has a *small diagonal* if, for any uncountable set $A \subset (X \times X) \setminus \Delta$ there exists an uncountable $B \subset A$ such that $\overline{B} \cap \Delta = \emptyset$. The *spread* $s(X)$ of a space X is the supremum of cardinalities of discrete subspaces of X and $ext(X) = \sup\{|D| : D \text{ is a closed and discrete subset of } X\}$. Now, $hl(X) = \sup\{l(Y) : Y \subset X\}$ is the hereditary Lindelöf number of X . The cardinal $iw(X) = \min\{\kappa : \text{the space } X \text{ has a weaker topology of weight } \kappa\}$ is called *i -weight* of X . Recall that $iw(X) \leq nw(X)$ and $hl(X) \leq nw(X)$ for any space X .

If X is a space and $A \subset X$ we say that a family \mathcal{B} of subsets of X is an outer network (base) of the set A in X if $(\mathcal{B} \subset \tau(X))$ and for any $U \in \tau(A, X)$ there exists $B \in \mathcal{B}$ such that $A \subset B \subset U$. Given an infinite cardinal κ , recall that $t(X) \leq \kappa$ if $\overline{A} = \bigcup\{\overline{B} : B \subset A \text{ and } |B| \leq \kappa\}$ for any $A \subset X$. A continuous map $f : X \rightarrow Y$ is *compact-covering* if for any $L \in \mathcal{K}(Y)$ there exists $K \in \mathcal{K}(X)$ such that $f(K) = L$. For any spaces X and Y the space $C_p(X, Y)$ consists of continuous functions from X to Y with the topology induced from Y^X . The space $C_p(X, \mathbb{R})$ is denoted by $C_p(X)$.

The rest of our notation is standard and follows [En]; our reference book on C_p -theory is [Ar2].

2. General properties of spaces dominated by second countable ones.

Our purpose is to find interesting classes in which domination by a second countable space coincides with the Lindelöf Σ -property. We show that this coincidence takes place for the spaces $C_p(X)$ and sometimes for the complements of the

diagonal of compact spaces. The following result summarizes the simplest properties of spaces dominated by second countable ones.

- 2.1. Theorem.** (a) Every Lindelöf Σ -space is dominated by a second countable space;
- (b) if X is dominated by a second countable space then any continuous image of X is also dominated by a second countable space;
- (c) if X is dominated by a second countable space then any closed subspace of X is also dominated by a second countable space;
- (d) if $X = \bigcup_{i \in \omega} X_i$ and X_i is dominated by a second countable space for all $i \in \omega$ then X is dominated by a second countable space;
- (e) if X_i is dominated by a second countable space for all $i \in \omega$ then the space $X = \prod_{i \in \omega} X_i$ is dominated by a second countable space;
- (f) if X is a space and $Y_i \subset X$ is dominated by a second countable space for all $i \in \omega$ then $Y = \bigcap_{i \in \omega} Y_i$ is also dominated by a second countable space;
- (g) a space X is Lindelöf Σ if and only if it is Dieudonné complete (i.e., homeomorphic to a closed subspace of a product of metrizable spaces) and dominated by a second countable space;
- (h) if X is dominated by a second countable space then $\text{ext}(X) = \omega$.

Proof. The statement of (a) was proved in the first paragraph of Introduction; the proofs of (b) and (c) are straightforward and can be left to the reader. To see that (d) is true suppose that X_i has an M_i -ordered compact cover $\mathcal{F}_i = \{P(K, i) : K \in \mathcal{K}(M_i)\}$ for some second countable space M_i for every $i \in \omega$. The space $M = \bigoplus_{i \in \omega} M_i$ is second countable; we identify every M_i with the corresponding clopen subset of M . Given any $K \in \mathcal{K}(M)$ the set $N_K = \{i \in \omega : K \cap M_i \neq \emptyset\}$ is finite so the set $F_K = \bigcup\{P(K \cap M_i, i) : i \in N_K\}$ is compact. It is immediate that the family $\{F_K : K \in \mathcal{K}(M)\}$ is an M -ordered compact cover of X .

(e) For each $i \in \omega$ fix a second countable space M_i and an M_i -ordered compact cover $\mathcal{F}_i = \{Q(K, i) : K \in \mathcal{K}(X_i)\}$ of the space X_i . For the space $M = \prod_{i \in \omega} M_i$ let $p_i : M \rightarrow M_i$ be the natural projection for every $i \in \omega$. Given any $K \in \mathcal{K}(M)$, the set $F_K = \prod\{Q(p_i(K), i) : i \in \omega\}$ belongs to $\mathcal{K}(X)$. It is an easy exercise that the family $\{F_K : K \in \mathcal{K}(M)\}$ is an M -ordered compact cover of X .

It is standard to deduce (f) from (c) and (e); the statement of (g) was proved in [CO2]. If X is dominated by a second countable space and D is a closed discrete subspace of X then D is also dominated by a second countable space by (c). Since D is also Dieudonné complete, it must be Lindelöf and hence countable by (g). This shows that $\text{ext}(X) = \omega$, i.e., (h) is proved. \square

2.2. Proposition. *The following conditions are equivalent for any space X :*

- (a) X has a \mathbb{P} -directed compact cover, i.e., X is dominated by the irrationals in the sense of [Tk];
- (b) X is \mathbb{P} -dominated;
- (c) X is dominated by a Polish space.

Proof. (a) \implies (b) Fix a \mathbb{P} -directed compact cover $\{Q(p) : p \in \mathbb{P}\}$ of the space X and let $\pi_i : \mathbb{P} \rightarrow \omega$ be the projection of \mathbb{P} onto its i -th factor, i.e., $\pi_i(s) = s(i)$ for any $s \in \mathbb{P}$. If $K \in \mathcal{K}(\mathbb{P})$ then $\pi_i(K)$ is a finite set so the number $s_K(i) = \max(\pi_i(K))$ is well-defined for any $i \in \omega$ and hence we have an element $s_K \in \mathbb{P}$ for any $K \in \mathcal{K}(\mathbb{P})$. It is immediate that $K \subset L$ implies that $s_K \leq s_L$; let $F_K = Q(s_K)$ for any $K \in \mathcal{K}(\mathbb{P})$. It is straightforward that $\mathcal{F} = \{F_K : K \in \mathcal{K}(\mathbb{P})\}$ is a \mathbb{P} -ordered family of compact subsets of X . To see that \mathcal{F} is a cover of X fix any point $x \in X$ and $p \in \mathbb{P}$ with $x \in Q(p)$. The set $K = \prod\{\{0, \dots, p(i)\} : i \in \omega\}$ is compact and $s_K = p$; as a consequence, $x \in Q(p) = Q(s_K) = F_K$ so \mathcal{F} is a compact \mathbb{P} -ordered cover of X , i.e., X is \mathbb{P} -dominated.

(b) \implies (a) Assume that the space X is \mathbb{P} -dominated and fix a respective compact cover $\{F_K : K \in \mathcal{K}(\mathbb{P})\}$. For any $p \in \mathbb{P}$ the set $K(p) = \prod\{\{0, \dots, p(i)\} : i \in \omega\}$ is compact; let $Q(p) = F_{K(p)}$. It is easy to see that the family $\mathcal{Q} = \{Q(p) : p \in \mathbb{P}\}$ is \mathbb{P} -directed. To see that \mathcal{Q} is a cover of X take a point $x \in X$; there exists $K \in \mathcal{K}(\mathbb{P})$ with $x \in F_K$. Consider the point $p \in \mathbb{P}$ such that $p(i) = \max(\pi_i(K))$ for every $i \in \omega$. Then $K \subset K(p)$ and hence $x \in F_K \subset F_{K(p)}$ so \mathcal{Q} is a \mathbb{P} -directed compact cover of X .

The implication (b) \implies (c) being clear, assume that a space X is dominated by a Polish space M and take a respective M -ordered compact cover $\{F(L) : L \in \mathcal{K}(M)\}$. There exists an open continuous onto map $\varphi : \mathbb{P} \rightarrow M$; observe that the family $\mathcal{F} = \{F(\varphi(K)) : K \in \mathcal{K}(\mathbb{P})\}$ is \mathbb{P} -ordered. To see that \mathcal{F} covers X take any point $x \in X$ and a compact set $L \subset M$ such that $x \in F(L)$. Any open map between Polish spaces is inductively perfect and hence compact-covering (see e.g., [En, 5.5.8]) so there exists $K \in \mathcal{K}(\mathbb{P})$ such that $\varphi(K) = L$. Therefore $x \in F(\varphi(K)) \in \mathcal{F}$ and hence \mathcal{F} is a \mathbb{P} -ordered compact cover of X , i.e., we settled (c) \implies (b). \square

2.3. Corollary. *A Dieudonné complete space is K -analytic if and only if it is dominated by a Polish space.*

Proof. It was proved in [CO2] that a Dieudonné complete space is K -analytic if and only if it has a \mathbb{P} -directed compact cover; Proposition 2.2 does the rest. \square

2.4. Corollary. *For any space X , if $C_p(X)$ is dominated by a Polish space then it is K -analytic.*

Proof. It was proved in [Tk] that any \mathbb{P} -dominated space $C_p(X)$ is K -analytic so we can apply Proposition 2.2 to finish our proof. \square

Cascales and Orihuela proved (using a different terminology), that if X is compact and $X^2 \setminus \Delta$ is strongly dominated by the irrationals then X is metrizable (see [CO1, Theorem 1]). It is a very interesting question whether the word “strongly” can be omitted in this statement. Our plan is to show that this is true under $\text{MA}(\omega_1)$. We will use the methods developed in [CKS] adapted to our situation. For the reader’s convenience we avoid citing very general technical results from [CKS] and give direct short proofs here for some particular cases we need.

2.5. Proposition. Suppose that X is dominated by a second countable space M and a collection $\{F_K : K \in \mathcal{K}(M)\}$ witnesses this. Take a countable base \mathcal{B} in M such that the union and the intersection of any finite subfamily of \mathcal{B} belongs to \mathcal{B} . For any $U \in \mathcal{B}$ let $G(U) = \bigcup\{F_K : K \in \mathcal{K}(M) \text{ and } K \subset U\}$. Fix a set $K \in \mathcal{K}(M)$ and a family $\mathcal{B}_K = \{U_n : n \in \omega\} \subset \mathcal{B}$ such that $U_{n+1} \subset U_n$ for each $n \in \omega$ and \mathcal{B}_K is an outer base of K in M ; then $F_K \subset C_K = \bigcap\{G(U) : U \in \mathcal{B}_K\}$. If $S = \{y_n : n \in \omega\} \subset X$ is a sequence such that $y_n \in G(U_n)$ for all $n \in \omega$, then

- (a) the set \overline{S} is compact and hence the set D of cluster points of S is non-empty;
- (b) there exists a compact set Q_K such that $D \subset Q_K \subset C_K$.

Proof. Take a set $K_n \in \mathcal{K}(M)$ such that $K_n \subset U_n$ and $y_n \in F_{K_n}$ for any $n \in \omega$. It is straightforward that the set $L_m = K \cup (\bigcup\{K_i : i \geq m\})$ is compact for any $m \in \omega$. The sequence $\{y_n\}$ is eventually in the compact set F_{L_m} which shows that the set \overline{S} is compact, $D \neq \emptyset$ and $D \subset F_{L_m}$ for any $m \in \omega$. Therefore D is contained in the compact set $Q_K = \bigcap\{F_{L_m} : m \in \omega\} \subset C_K$ as promised. \square

2.6. Proposition. Suppose that X is dominated by a second countable space M and a collection $\{F_K : K \in \mathcal{K}(M)\}$ witnesses this. Fix a countable base \mathcal{B} in M such that the union and the intersection of any finite subfamily of \mathcal{B} belongs to \mathcal{B} . For any $U \in \mathcal{B}$ let $G(U) = \bigcup\{F_K : K \in \mathcal{K}(M) \text{ and } K \subset U\}$. Then there exists a family \mathcal{C} in the space X with the following properties:

- (a) if $C \in \mathcal{C}$ and $A \subset C$ is a countable set then the set \overline{A} is compact and $\overline{A} \subset C$; in particular, each $C \in \mathcal{C}$ is countably compact;
- (b) for every $K \in \mathcal{K}(M)$ there exists a set $C_K \in \mathcal{C}$ such that $F_K \subset C_K$ and hence \mathcal{C} is a cover of X ;
- (c) the family $\mathcal{N} = \{G(U) : U \in \mathcal{B}\}$ is a network with respect to \mathcal{C} .

Proof. Fix any compact subset K of the space M and observe that we can choose a family $\mathcal{B}_K = \{U_n : n \in \omega\} \subset \mathcal{B}$ such that $U_{n+1} \subset U_n$ for each $n \in \omega$ and \mathcal{B}_K is an outer base of K in M . It is evident that $F_K \subset C_K = \bigcap\{G(U) : U \in \mathcal{B}_K\}$. Let $\mathcal{C} = \{C_K : K \in \mathcal{K}(M)\}$; it is clear that the property (b) holds for \mathcal{C}_K .

If $K \in \mathcal{K}(M)$ and $\{G(U_n) : n \in \omega\}$ is not a network for C_K then we can choose a point $y_n \in G(U_n) \setminus W$ for some $W \in \tau(C_K, X)$. The sequence $\{y_n\}$ must have a cluster point in C_K by Proposition 2.5 which contradicts the fact that $\{y_n\} \subset X \setminus W$ while $C_K \subset W$. Therefore the family \mathcal{C} has the property (c).

Furthermore, if $A \subset C_K$ is countable then we can choose an enumeration $\{y_n : n \in \omega\}$ of the set A . It is clear that $y_n \in G(U_n)$ for all $n \in \omega$ and hence we can apply Proposition 2.5 again to see that $\overline{A} = \overline{\{y_n : n \in \omega\}}$ is compact. If $x \in \overline{A} \setminus C_K$ then $x \in \overline{A} \setminus A$ and hence x is a cluster point of the sequence $S = \{y_n\}$. However, all cluster points of S belong to C_K by Proposition 2.5. This contradiction shows that $\overline{A} \subset C_K$ so (a) is proved as well. \square

Given a space X recall that a set $A \subset X$ is *relatively countably compact* if every sequence in A has a cluster point in X . The following result was implicitly proved in [CO2] and [CKS].

2.7. Corollary. *Suppose that, in a space X , every relatively countably compact set has compact closure. Then X is dominated by a second countable space if and only if it has the Lindelöf Σ -property. In particular, an angelic space X is dominated by a second countable space if and only if X is Lindelöf Σ .*

Proof. It suffices to prove necessity so assume that X is dominated by a second countable space. It follows from Proposition 2.6 that we can find a cover \mathcal{C} of the space X such that every $C \in \mathcal{C}$ is countably compact and there exists a countable network \mathcal{N} with respect to \mathcal{C} . The family $\mathcal{F} = \{\overline{C} : C \in \mathcal{C}\}$ is a cover of X and all elements of \mathcal{F} are compact. It is standard that $\mathcal{M} = \{\overline{N} : N \in \mathcal{N}\}$ is a countable network with respect to \mathcal{F} so X is a Lindelöf Σ -space. \square

2.8. Theorem. *Suppose that Z is a compact space of countable tightness. Then a set $X \subset Z$ is dominated by a second countable space if and only if X has the Lindelöf Σ -property.*

Proof. Fix any set $X \subset Z$ and assume that X is dominated by a second countable space. For any set $A \subset X$ we denote by $\text{cl}_X(A)$ (or $\text{cl}_Z(A)$) the closure of the set A in the space X (or in Z respectively). By Proposition 2.6, there exists a cover \mathcal{C} of the space X and a countable network \mathcal{N} with respect to \mathcal{C} such that for every $C \in \mathcal{C}$ and any countable $A \subset C$ the set $\text{cl}_X(A)$ is compact and contained in C .

If $C \in \mathcal{C}$ and C is not closed in Z then we can find a point $x \in \text{cl}_Z(C) \setminus C$. By countable tightness of Z , there exists a countable $A \subset C$ such that $x \in \text{cl}_Z(A)$. The set $F = \text{cl}_X(A) \subset C$ is compact and hence closed in Z ; as a consequence, $x \in \text{cl}_Z(A) \subset F \subset C$. This contradiction shows that every $C \in \mathcal{C}$ is compact being closed in X . Thus \mathcal{N} is a countable network with respect to the compact cover \mathcal{C} of the space X , i.e., X has the Lindelöf Σ -property. \square

2.9. Theorem. *If X is a compact space with $t(X) \leq \omega$ and $X^2 \setminus \Delta$ is dominated by a second countable space then X is metrizable.*

Proof. The space X^2 also has countable tightness [Ar1, Theorem 2.3.3] so we can apply Theorem 2.8 to the set $X^2 \setminus \Delta \subset X \times X$ to conclude that $X^2 \setminus \Delta$ is a Lindelöf Σ -space; this easily implies that the diagonal Δ is a G_δ -subset of $X \times X$ and hence X is metrizable by [En, 3.12.22(e)]. \square

2.10. Corollary. *If X is a Corson compact space or a first countable compact space such that $X^2 \setminus \Delta$ is dominated by a second countable space then X is metrizable.*

2.11. Theorem. *If X is a dyadic compact space and $X^2 \setminus \Delta$ is dominated by a second countable space then X is metrizable.*

Proof. If X is first countable then it is metrizable by [En, 3.12.12(e)]. Therefore we can assume that there exists a point $x \in X$ of uncountable character in X . Apply [En, 3.12.12(i)] to find an uncountable one-point compactification A of a discrete space such that $A \subset X$ and x is the unique non-isolated point of A . Then

$B = (A \setminus \{x\}) \times \{x\}$ is an uncountable closed discrete subspace of $(X \times X) \setminus \Delta$ while we have $\text{ext}(X^2 \setminus \Delta) = \omega$ by Theorem 2.1(h), a contradiction. \square

The above results show that, to prove that any compact space X with $X^2 \setminus \Delta$ dominated by a second countable space is metrizable, it suffices to show that any such space has a countable tightness. While we don't know whether this implication is true in general, we do present some partial progress in this direction.

2.12. Theorem. *Assume $MA(\omega_1)$ and suppose that X is a compact space such that $X^2 \setminus \Delta$ is \mathbb{P} -dominated. Then X has a small diagonal and hence $t(X) = \omega$.*

Proof. Suppose that $A = \{z_\alpha : \alpha < \omega_1\} \subset X^2 \setminus \Delta$ and $\alpha \neq \beta$ implies $z_\alpha \neq z_\beta$. Fix a \mathbb{P} -directed cover $\{K_p : p \in \mathbb{P}\}$ of compact subsets of $X^2 \setminus \Delta$. Take $p_\alpha \in \mathbb{P}$ such that $z_\alpha \in K_{p_\alpha}$ for any $\alpha < \omega_1$.

It follows from $MA(\omega_1)$ that there exists $p \in \mathbb{P}$ such that $p_\alpha \leq^* p$ for any $\alpha < \omega_1$. The set $P = \bigcup \{K_q : q \in \mathbb{P} \text{ and } q =^* p\}$ is σ -compact and $A \subset P$. Consequently, there is $q \in \mathbb{P}$ for which $K_q \cap A$ is uncountable; therefore the set $K_q \cap A$ witnesses the small diagonal property of X . Since no space with a small diagonal can have a convergent ω_1 -sequence, it follows from [JuS, Theorem 1.2] that X has no free sequences of length ω_1 , i.e., $t(X) \leq \omega$. \square

2.13. Corollary. *Under $MA(\omega_1)$, if X is a compact space such that $X^2 \setminus \Delta$ is dominated by a Polish space then X is metrizable.*

Proof. Apply Proposition 2.2 to see that the space $X^2 \setminus \Delta$ is dominated by \mathbb{P} so $t(X) \leq \omega$ by Theorem 2.11 and hence X is metrizable by Theorem 2.9. \square

In the rest of this section we study the spaces hereditarily dominated by a second countable space. The motivation here is a result of Hodel established in [Ho, Corollary 4.13]; it says that any hereditarily Lindelöf Σ -space is cosmic. We will look at this hereditary property in function spaces to show that a somewhat stronger statement is true in a general situation under Martin's Axiom.

The following fact is an immediate consequence of [Tk, Proposition 2.7].

2.14. Proposition. *If X is a space which has a countable network modulo a cover of X by countably compact sets then $C_p(X)$ is Lindelöf Σ -framed, i.e., there is a Lindelöf Σ -space L such that $C_p(X) \subset L \subset \mathbb{R}^X$.*

2.15. Theorem. *A space $C_p(X)$ is dominated by a second countable space if and only if it is Lindelöf Σ .*

Proof. We must only prove necessity. Suppose that $C_p(X)$ is dominated by a second countable space M and fix a family $\{F_K : K \in \mathcal{K}(M)\}$ which witnesses this. It follows from Proposition 2.14 and Proposition 2.6 that $C_p(C_p(X))$ is Lindelöf Σ -framed. Applying [Ok, Theorem 3.5] we conclude that $v(C_p(X))$ is a Lindelöf Σ -space and hence vX is a Lindelöf Σ -space by [Ok, Corollary 3.6].

Let $\pi : C_p(vX) \rightarrow C_p(X)$ be the restriction map. If $G_K = \pi^{-1}(F_K)$ then G_K is compact for any $K \in \mathcal{K}(M)$ (see [Tk, Theorem 2.6]). It is clear that $\mathcal{G} = \{G_K : K \in \mathcal{K}(M)\}$ is a cover of $C_p(vX)$ which shows that $C_p(vX)$ is dominated by M . By Proposition 2.6 we can find a countable network \mathcal{N} modulo a cover \mathcal{C} of the space $C_p(vX)$ such that every $C \in \mathcal{C}$ is countably compact. Every countably compact subset of $C_p(vX)$ is compact by [Ar2, Proposition IV.9.10] (see also [Or]) so \mathcal{C} consists of compact subsets of $C_p(vX)$ and hence $C_p(vX)$ is a Lindelöf Σ -space. Therefore $C_p(X)$ is also Lindelöf Σ -space being a continuous image of $C_p(vX)$. \square

2.16. Lemma. *If every subspace of a space X is realcompact (i.e., X is hereditarily realcompact) and dominated by a second countable space then X is cosmic.*

Proof. Every subspace of X has to be Lindelöf Σ by Theorem 2.1(g) so we can apply [Ho, Corollary 4.13] to conclude that X is cosmic. \square

2.17. Theorem. *Under Martin's axiom, the following conditions are equivalent for any space X :*

- (a) every subspace of X is dominated by a second countable space;
- (b) the space X is cosmic.

Proof. Every subspace of a cosmic space is cosmic and hence Lindelöf Σ so it is dominated by a second countable space by Theorem 2.1(a). This proves that (b) \implies (a); observe that no additional axioms are needed for this conclusion.

Now assume that there exist non-cosmic spaces which are hereditarily dominated by a second countable space and call every such space a *counterexample*. Observe first that a counterexample cannot be hereditarily Lindelöf by Lemma 2.16. Therefore, if X is a counterexample then we can find a right-separated subspace $Y \subset X$ such that $|Y| = \omega_1$. It is immediate that Y is also a counterexample so we can assume, without loss of generality, that $X = Y$, i.e., X is a scattered space. If every countably compact subspace of X is compact and $Y \subset X$ then we can apply Proposition 2.6 to find a cover \mathcal{C} of Y by countably compact (and hence compact) subspaces such that there exists a countable network modulo \mathcal{C} . This proves that every $Y \subset X$ is Lindelöf Σ and hence X is cosmic by [Ho, Corollary 4.13], which is a contradiction.

Therefore we can find an uncountable countably compact subspace $Y \subset X$; it is clear that Y is also a counterexample. Thus we can assume, without loss of generality, that X is countably compact. It follows from Theorem 2.1(h) that $s(X) \leq \omega$ and hence X is hereditarily separable (see [Ju, 2.12]).

If Y is a subspace of X then let $I(Y)$ be the set of isolated points of Y ; if $Y \neq \emptyset$ then $I(Y) \neq \emptyset$ because the space X is scattered. Let $X_0 = X$; if α is a countable ordinal and we have X_α then $X_{\alpha+1} = X_\alpha \setminus I(X_\alpha)$. If α is a limit ordinal and we have X_β for every $\beta < \alpha$ then $X_\alpha = \bigcap_{\beta < \alpha} X_\beta$. This gives us a strictly decreasing ω_1 -sequence $\{X_\alpha : \alpha < \omega_1\}$ of closed subsets of X such that $X \setminus X_\alpha$ is countable and hence $X_\alpha \neq \emptyset$ for any $\alpha < \omega_1$.

The set $Y = \bigcup_{\alpha < \omega_1} (X \setminus X_\alpha)$ is a counterexample because it has cardinality ω_1 . The space Y is an increasing union of countable open subsets of X . Therefore every point of Y has a countable countably compact neighbourhood, i.e., Y is locally compact and locally countable. The one-point compactification of Y is an uncountable compact scattered hereditarily separable space. Such spaces do not exist under $\text{MA} + \neg\text{CH}$ (see [Ro, Theorem 6.4.1]) so if CH does not hold then our proof is over.

Finally, assume that CH holds and observe that Y is first countable so every countably compact subspace of Y is closed in Y . Therefore every countably compact subset of Y is uniquely determined by its countable dense subset and hence the family \mathcal{P} of uncountable countably compact subspace of Y has cardinality at most $\omega_1^\omega = \mathfrak{c} = \omega_1$.

It is standard that we can find disjoint subsets A, B of the space Y such that $Y = A \cup B$ and $A \cap P \neq \emptyset \neq B \cap P$ for any $P \in \mathcal{P}$. In particular, every countably compact subset of A as well as every countably compact subspace of B is countable and hence compact. This, together with Proposition 2.6 implies that both A and B are hereditarily Lindelöf Σ so we can apply [Ho, Corollary 4.13] again to see that $nw(A) = nw(B) = \omega$ and hence $Y = A \cup B$ is cosmic which is a contradiction. \square

If a space $C_p(X)$ is hereditarily dominated by a second countable space then no additional axioms are needed to obtain the same conclusion as in Theorem 2.17.

2.18. Proposition. *If every subspace of a space $C_p(X)$ is dominated by a second countable space then $C_p(X)$ is cosmic.*

Proof. We have $s(C_p(X)) = \omega$ by Theorem 2.1(h); besides, $C_p(X)$ is a Lindelöf Σ -space by Theorem 2.15. If $C_p(X)$ is not hereditarily Lindelöf then we can find an uncountable right-separated subspace $Y \subset C_p(X)$ (see [Ju, Theorem 2.9(b)]). Every right-separated space of countable spread must be hereditarily separable (see [Ju, Theorem 2.12]) so Y is separable. In the space $C_p(X)$ the closure of every countable subset is cosmic by [Ar3, Theorem 7.21] so we can conclude that $nw(Y) \leq \omega$ and, in particular, $hl(Y) \leq \omega$ which is a contradiction. This proves that $C_p(X)$ is hereditarily Lindelöf so it follows from Lemma 2.16 that $C_p(X)$ is cosmic. \square

3. Strong domination by second countable spaces.

Say that a space X is *strongly dominated by a space M* if there exists a cover $\mathcal{F} = \{F_K : K \in \mathcal{K}(M)\}$ of the space X such that F_K is compact for any $K \in \mathcal{K}(M)$ and the family \mathcal{F} swallows all compact subsets of X in the sense that for any compact $C \subset X$ there is $K \in \mathcal{K}(M)$ such that $C \subset F_K$.

The following two results seem to be a good motivation for a systematic study of the class \mathcal{M}^* of spaces which are strongly dominated by second countable ones.

3.1. Theorem (Christensen, [Chr, Theorem 3.3]). *A second countable space is strongly \mathbb{P} -dominated if and only if it is completely metrizable.*

3.2. Theorem (Cascales and Orihuela, [CO1, Theorem 1]). *If X is a compact space such that $(X \times X) \setminus \Delta$ is strongly \mathbb{P} -dominated then X is metrizable. Here $\Delta = \{(x, x) : x \in X\}$ is the diagonal of the space X .*

3.3. Proposition. (a) *If X is strongly dominated by a second countable space and Y is a compact-covering image of X then Y is strongly dominated by a second countable space;*

(b) *every \aleph_0 -space is strongly dominated by a second countable space;*

(c) *if X is strongly dominated by a second countable space then every closed subspace of X is also strongly dominated by a second countable space;*

(d) *if X_i is strongly dominated by a second countable space for every $i \in \omega$ then $\prod_{i \in \omega} X_i$ is strongly dominated by a second countable space;*

(e) *if X is a space and $Y_i \subset X$ is strongly dominated by a second countable space for each $i \in \omega$ then $Y = \bigcap_{i \in \omega} Y_i$ is also strongly dominated by a second countable space.*

Proof. Suppose that X is strongly dominated by a second countable space M and $f : X \rightarrow Y$ is a compact-covering map. Let $\{F_K : K \in \mathcal{K}(M)\}$ be the family which witnesses that X is strongly dominated by M and consider the family $\mathcal{F} = \{f(F_K) : K \in \mathcal{K}(M)\}$. It is clear that \mathcal{F} consists of compact subsets of Y and $K \subset L$ implies $f(F_K) \subset f(F_L)$. If P is a compact subset of Y then there exists a compact subset $Q \subset X$ such that $f(Q) = P$. Pick a set $K \in \mathcal{K}(M)$ such that $Q \subset F_K$ and observe that $P = f(Q) \subset f(F_K)$. Therefore the family \mathcal{F} witnesses that Y is strongly dominated by M , i.e., we proved (a).

The item (b) follows from (a) and the fact that every \aleph_0 -space is a compact-covering image of a second countable space [Mi, Theorem 11.4]. The proof of (c) is straightforward and can be left to the reader.

Next assume that X_i is strongly dominated by a second countable space M_i and fix a respective family $\mathcal{F}_i = \{F_i(K) : K \in \mathcal{K}(M_i)\}$ for any $i \in \omega$. The space $M = \prod_{i \in \omega} M_i$ is second countable; let $\pi_i : M \rightarrow M_i$ be the natural projection for each $i \in \omega$. If $K \in \mathcal{K}(M)$ then $F_K = \prod_{i \in \omega} F_i(\pi_i(K))$ is easily seen to be a compact subset of $X = \prod_{i \in \omega} X_i$. Let $p_i : X \rightarrow X_i$ be the natural projection for every $i \in \omega$.

The family $\mathcal{F} = \{F_K : K \in \mathcal{K}(M)\}$ witnesses that X is strongly dominated by M . Indeed, if Q is a compact subset of X then we can choose $K_i \in \mathcal{K}(M_i)$ such that $p_i(Q) \subset F_i(K_i)$ for each $i \in \omega$; for the set $K = \prod_{i \in \omega} K_i$ we have $Q \subset F_K$. It is immediate that $K \subset L$ implies $F_K \subset F_L$ so we settled (d). As to (e), observe that Y is homeomorphic to a closed subspace of $\prod_{i \in \omega} Y_i$ so we can apply (c) and (d) to finish the proof. \square

3.4. Proposition. *The space ω_1 with its interval topology is strongly dominated by the space of rational numbers.*

Proof. Given a compact set $K \subset \mathbb{Q}$, let $\alpha_K \in \omega_1$ be the minimal ordinal such that $F_K = \{\beta : \beta < \alpha_K\}$, as a subspace of ω_1 , is homeomorphic to K . Such an ordinal

α_K exists by [MS, Theorem 1]. It is clear that the family $\mathcal{F} = \{F_K : K \in \mathcal{K}(\mathbb{Q})\}$ is \mathbb{Q} -ordered.

Suppose that L is a compact subset of ω_1 and choose an ordinal $\alpha < \omega_1$ such that $L \subset \{\beta : \beta < \alpha\}$. It is easy to see that there exists a countable ordinal $\gamma > \alpha$ such that $Q = \{\beta : \beta < \gamma\}$ is a compact subset of ω_1 and no initial segment of Q is homeomorphic to Q . The space \mathbb{Q} is also universal for all countable compact spaces so there exists $K \subset \mathbb{Q}$ with $K \simeq Q$. It is clear that $\alpha_K = \gamma$ and hence $L \subset \{\beta : \beta < \alpha\} \subset Q = F_K$. This shows that \mathcal{F} is a \mathbb{Q} -ordered compact cover of ω_1 which swallows all compact subsets of ω_1 , i.e., ω_1 is strongly \mathbb{Q} -dominated. \square

3.5. Corollary. *Under $MA+\neg CH$ there exists a strongly \mathbb{Q} -dominated space which is not \mathbb{P} -dominated.*

Proof. The space ω_1 is not \mathbb{P} -dominated under $MA+\neg CH$ (see [Tk, Theorem 3.6]) so apply Proposition 3.4 to see that ω_1 is as promised. \square

Proposition 3.3(b) and Proposition 3.4 show that \mathcal{M}^* is strictly larger than the class of \aleph_0 -spaces. Therefore it is natural to ask when domination by a second countable space must imply the \aleph_0 -property. Recall that a space is called *submetrizable* if it has a weaker metrizable topology.

3.6. Theorem. *The following conditions are equivalent for any space X :*

- (a) X is an \aleph_0 -space;
- (b) X is strongly dominated by a second countable space and $iw(X) \leq \omega$;
- (c) X is submetrizable and strongly dominated by a second countable space.

Proof. Every \aleph_0 -space X is cosmic and hence $iw(X) \leq \omega$; this, together with Proposition 3.3(b), shows that (a) \implies (b). The implication (b) \implies (c) being trivial assume that X is submetrizable and strongly dominated by a second countable space. It follows from [CO2, Theorem 4] that X is a Lindelöf Σ -space so its weaker metrizable topology must be second countable, i.e., $iw(X) \leq \omega$.

Fix an M -ordered family $\{F_K : K \in \mathcal{K}(M)\}$ of compact subsets of X such that every $L \in \mathcal{K}(X)$ is contained in some F_K . Apply Proposition 2.6 to find a family \mathcal{C} of countably compact (and hence compact) subsets of X such that some countable family \mathcal{N} is a network modulo \mathcal{C} and, for every $K \in \mathcal{K}(M)$ there exists $C_K \in \mathcal{C}$ such that $F_K \subset C_K$. In particular, the family \mathcal{C} swallows all compact subsets of X .

Taking the closures of the elements of \mathcal{N} we will still have a network modulo \mathcal{C} so we can assume, without loss of generality, that \mathcal{N} consists of closed subsets of X . Fix a second countable topology μ on the set X such that $\mu \subset \tau(X)$. The space (X, μ) has a countable closed network \mathcal{P} modulo all compact subsets of (X, μ) . Observe that the identity map $id : X \rightarrow (X, \mu)$ is continuous and hence any compact subset of X is also compact in (X, μ) . Consider the family \mathcal{Q} of all finite unions and finite intersections of the elements of the family $\mathcal{P} \cup \mathcal{N}$; we claim that \mathcal{Q} is a network for all compact subsets of X .

Indeed, take any $L \in \mathcal{K}(X)$ and $U \in \tau(L, X)$. There exists $C \in \mathcal{C}$ such that $L \subset C$. The set $C \setminus U$ does not meet L so there exists $P \in \mathcal{P}$ such that $L \subset P$ and $P \cap (C \setminus U) = \emptyset$. The set $P' = P \setminus U$ does not meet C so we can find a set $N \in \mathcal{N}$ such that $C \subset N \subset X \setminus P'$. The set $Q = N \cap P$ belongs to \mathcal{Q} and $L \subset Q \subset U$ so the family \mathcal{Q} witnesses that X is an \aleph_0 -space. \square

3.7. Remark. *Adapting to our situation the proof of the implication (ii) \implies (i) in Theorem 6 of [CO2] gives another direct (and somewhat shorter) way to establish the implication (c) \implies (a) in Theorem 3.6.*

3.8. Corollary. *Under Martin's Axiom, every subspace of a space X is strongly dominated by a second countable space if and only if X is an \aleph_0 -space.*

Proof. If X is an \aleph_0 -space then every subspace of X is also \aleph_0 -space so X is hereditarily strongly dominated by a second countable space by Proposition 3.3(b); this proves sufficiency.

If X is hereditarily strongly dominated by a second countable space then we can apply Theorem 2.17 to convince ourselves that X is cosmic and hence $iw(X) \leq \omega$. Now it follows from Theorem 3.6 that X is an \aleph_0 -space. \square

Given an infinite cardinal κ say that a space X is κ -hemicompact if there exists a family \mathcal{F} of compact subsets of X such that $|\mathcal{F}| \leq \kappa$ and \mathcal{F} swallows all compact subsets of X , i.e., for any $K \in \mathcal{K}(X)$ there exists $F \in \mathcal{F}$ such that $K \subset F$. Observe that a space is hemicompact if and only if it is ω -hemicompact.

3.9. Theorem. *The σ -product $S_\kappa = \{x \in \mathbb{D}^\kappa : |x^{-1}(1)| < \omega\}$ of the space \mathbb{D}^κ is not κ -hemicompact for any infinite cardinal κ .*

Proof. Denote by u the point of \mathbb{D}^κ which is identically zero on κ and hence $u^{-1}(1) = \emptyset$. Take any family $\mathcal{F} = \{F_\alpha : \alpha < \kappa\}$ of compact subsets of S_κ . The set S_κ is not compact so we can pick a point $x_0 \in S_\kappa \setminus F_0$. Proceeding inductively assume that $\alpha < \kappa$ and we have chosen a set $\{x_\beta : \beta < \alpha\}$ with the following properties:

- (1) $x_\beta \in S_\kappa \setminus F_\beta$ for any $\beta < \alpha$;
- (2) the family $\{x_\beta^{-1}(1) : \beta < \alpha\}$ is disjoint.

Observe that the set $A = \bigcup \{x_\beta^{-1}(1) : \beta < \alpha\}$ has cardinality strictly less than κ . Therefore the subspace $Y = \{x \in S_\kappa : x(A) = 0\}$ is not compact so we can choose a point $x_\alpha \in Y \setminus F_\alpha$; it is immediate that the conditions (1) and (2) are still satisfied for the set $\{x_\beta : \beta \leq \alpha\}$. Thus we can construct a set $\{x_\alpha : \alpha < \kappa\}$ for which the properties (1) and (2) hold for any $\alpha < \kappa$.

It follows from (2) that the set $K = \{x_\beta : \beta < \kappa\} \cup \{u\}$ is compact; the property (1) shows that $x_\beta \in K \setminus F_\beta$ for any $\beta < \kappa$ and therefore no element of the family \mathcal{F} swallows the set K . \square

3.10. Theorem. *Under the Continuum Hypothesis (CH) if a space X is compact and $C_p(X)$ is strongly dominated by a second countable space then X is countable and hence $C_p(X)$ is second countable.*

Proof. Apply Theorem 2.15 to see that $C_p(X)$ is a Lindelöf Σ -space and hence X is Gul'ko compact. If the space X is not scattered then we can find a countable dense-in-itself set $A \subset X$. The space $K = \overline{A}$ is compact, second countable and metrizable [Ar3, Theorem 7.21] so $C_p(K)$ embeds in $C_p(X)$ as a closed subspace [Ar3, Theorem 4.1]. This implies, by Proposition 3.3(c), that $C_p(K)$ is strongly dominated by a second countable space. Since $iw(C_p(K)) \leq nw(C_p(K)) = \omega$, we can apply Theorem 3.6 to convince ourselves that $C_p(K)$ is an \aleph_0 -space so K is countable by [Mi, Proposition 10.7]. However, K has no isolated points; this contradiction shows that X has to be scattered.

The set D of isolated points of the space X is dense in X ; if D is countable then X is second countable so we can apply Theorem 3.6 again to see that $C_p(X)$ is an \aleph_0 -space and hence X is countable by [Mi, Proposition 10.7]. Therefore we can assume that $\kappa = |D| \geq \omega_1$; consider the space Y which is obtained from X by contracting the set $F = X \setminus D$ to a point. It is evident that Y is a compact space with a unique non-isolated point, i.e., Y is homeomorphic to the one-point compactification A_κ of a discrete space of cardinality κ . The space Y is a continuous closed image of X so $C_p(Y)$ is homeomorphic to a closed subspace of $C_p(X)$. Thus $C_p(Y) \simeq C_p(A_\kappa)$ is strongly dominated by a second countable space.

It is an easy exercise that the space $C_p(A_\kappa)$ is homeomorphic to the Σ_* -product $\Omega = \{x \in \mathbb{R}^\kappa : \text{the set } \{\alpha < \kappa : |x(\alpha)| \geq \varepsilon\} \text{ is finite for any } \varepsilon > 0\}$ of the space \mathbb{R}^κ . Furthermore, $\Omega \cap \mathbb{D}^\kappa = S_\kappa = \{x \in \mathbb{D}^\kappa : x^{-1}(1) \text{ is finite}\}$ so S_κ is a closed subset of Ω ; in particular, S_κ is strongly dominated by a second countable space M . Let $\mathcal{F} = \{F_K : K \in \mathcal{K}(M)\}$ be a family of compact subsets of S_κ which witnesses this. However, $|\mathcal{K}(M)| \leq \mathfrak{c} = \omega_1$ so $|\mathcal{F}| \leq \omega_1$ and hence S_κ is ω_1 -hemicompact; since $\kappa \geq \omega_1$, we have obtained a contradiction with Theorem 3.9. \square

It is not difficult to deduce the following theorem from a general result proved by M. Muñoz in her PhD thesis (see [Mu, Theorem 2.10.1]). This result was also published in [CMO, Proposition 5.1]. For the reader's convenience we chose to avoid dealing with uniformities and give a direct topological proof here.

3.11. Theorem. *A compact space X is metrizable if and only if $X^2 \setminus \Delta$ is strongly dominated by a second countable space.*

Proof. The necessity being evident fix a second countable space E and a family $\mathcal{F} = \{F(Q) : Q \in \mathcal{K}(E)\}$ of compact subsets of $X^2 \setminus \Delta$ which witnesses that $X^2 \setminus \Delta$ is strongly E -dominated. Denote by C the subspace $C_p(X, [0, 1])$ of the space $C_p(X)$ and let $I = [0, 1]$. For the space $M = E^\mathbb{N}$ let $\pi_n : M \rightarrow E$ be the natural projection onto the n -th factor of M .

For every $K \in \mathcal{K}(M)$ consider the set $H_n = \{f \in I^X : |f(x) - f(y)| \leq \frac{1}{n} \text{ for any } (x, y) \in X^2 \setminus F(\pi_n(K))\}$ for each $n \in \mathbb{N}$ and let $G_K = \bigcap \{H_n : n \in \mathbb{N}\}$. It is

immediate that $K \subset L$ implies $G_K \subset G_L$ for any $K, L \in \mathcal{K}(M)$. We omit a simple proof of the fact that the set G_K is closed in I^X and hence compact. To see that $G_K \subset C$ take any $f \in G_K$, $x \in X$ and $\varepsilon > 0$. If $n \in \mathbb{N}$ and $\frac{1}{n} < \varepsilon$ then the set $U = \{y \in X : (x, y) \notin F(\pi_n(K))\}$ is an open neighbourhood of x in X and we have the inclusions

$$f(U) \subset [f(x) - 1/n, f(x) + 1/n] \subset (f(x) - \varepsilon, f(x) + \varepsilon)$$

which show that f is continuous at the point x . Thus G_K is a compact subset of C for any $K \in \mathcal{K}(M)$.

To see that $\mathcal{G} = \{G_K : K \in \mathcal{K}(M)\}$ is a cover of C take any $f \in C$. Then $O_n = \{(x, y) \in X^2 : |f(x) - f(y)| < 1/n\}$ is an open neighbourhood of Δ so the set $P_n = X^2 \setminus O_n \subset X^2 \setminus \Delta$ is compact for any $n \in \mathbb{N}$. The family \mathcal{F} swallows all compact subsets of $X^2 \setminus \Delta$ and hence we can find a set $K_n \in \mathcal{K}(M)$ such that $P_n \subset F(K_n)$ for all $n \in \mathbb{N}$. It is straightforward that $f \in G_K$ for the compact set $K = \prod\{K_n : n \in \mathbb{N}\}$ of the space M .

This proves that C is dominated by M ; since countably compact subsets of C are compact, we can apply Proposition 2.6 to see that there exists a countable network modulo a compact cover of C , i.e., the space C is Lindelöf Σ . The space X being compact, $C_p(X)$ is also Lindelöf Σ being the countable union of subspaces homeomorphic to C . It is easy to see that the space X^2 embeds in $C_p(C_p(X))$ whence $l(X^2 \setminus \Delta) = ext(X^2 \setminus \Delta) = \omega$ (see Theorem 2.1(h) and [Bat, Theorem 1']). Therefore $X^2 \setminus \Delta$ is Lindelöf; this easily implies that Δ is a G_δ -subset of $X \times X$ so X is metrizable by [En, 4.2.B]. \square

3.12. Corollary. *Suppose that X is a compact space, M is a second countable space and we have a family $\mathcal{G} = \{U_K : K \in \mathcal{K}(M)\}$ of neighbourhoods of the diagonal Δ in the space $X \times X$ such that $U_K \subset U_L$ whenever $L \subset K$. If, additionally, $\bigcap\{\bar{G} : G \in \mathcal{G} = \Delta\}$ then X is metrizable.*

Proof. Let $F_K = (X \times X) \setminus \text{Int}(U_K)$ for any $K \in \mathcal{K}(M)$. It is immediate that $F_K \subset F_L$ if $K \subset L$, i.e., the family $\mathcal{F} = \{F_K : K \in \mathcal{K}(M)\}$ is ordered by M . The equality $\bigcap\{\bar{G} : G \in \mathcal{G}\} = \Delta$ shows that $\bigcup\{\text{Int}(F_K) : K \in \mathcal{K}(M)\} = X^2 \setminus \Delta$. Given a compact set $F \subset X^2 \setminus \Delta$, the family $\{\text{Int}(F_K) : K \in \mathcal{K}(M)\}$ is an open cover of F so we can find $K_1, \dots, K_n \in \mathcal{K}(M)$ such that $F \subset \text{Int}(F_{K_1}) \cup \dots \cup \text{Int}(F_{K_n}) \subset F_K$ for $K = K_1 \cup \dots \cup K_n \in \mathcal{K}(M)$. Therefore the family \mathcal{F} witnesses that $X^2 \setminus \Delta$ is strongly dominated by the second countable space M and hence X is metrizable by Theorem 3.11. \square

3. Open problems.

One of the niceties of the concept of domination by a second countable space is a possibility to obtain new metrization theorems for compact spaces. We already saw that if X compact and $(X \times X) \setminus \Delta$ is strongly dominated by a second countable space then X is metrizable. The most interesting question here is whether we can omit the word “strongly” in the above statement.

- 3.1. Problem.** *Let X be a compact space such that $X^2 \setminus \Delta$ is \mathbb{P} -dominated. Is it true in ZFC that X must be metrizable?*
- 3.2. Problem.** *Let X be a compact space such that $X^2 \setminus \Delta$ is \mathbb{Q} -dominated. Is it true in ZFC that X must be metrizable?*
- 3.3. Problem.** *Let X be a compact space such that $X^2 \setminus \Delta$ is M -dominated for some separable metrizable space M . Is it true in ZFC that X must be metrizable?*
- 3.4. Problem.** *Suppose that X is a K -analytic space such that $X^2 \setminus \Delta$ is strongly \mathbb{P} -dominated. Must X be cosmic?*
- 3.5. Problem.** *Let X be a K -analytic space such that $X^2 \setminus \Delta$ is \mathbb{P} -dominated. Must X be cosmic?*
- 3.6. Problem.** *Suppose that X is a Lindelöf Σ -space such that $X^2 \setminus \Delta$ is strongly \mathbb{P} -dominated. Must X be cosmic?*
- 3.7. Problem.** *Let X be a Lindelöf Σ -space such that $X^2 \setminus \Delta$ is \mathbb{P} -dominated. Must X be cosmic?*
- 3.8. Problem.** *Let X be a Lindelöf Σ -space such that $X^2 \setminus \Delta$ is \mathbb{Q} -dominated. Must X be cosmic?*
- 3.9. Problem.** *Suppose that $C_p(X)$ is strongly \mathbb{Q} -dominated. Must the space X be countable?*
- 3.10. Problem.** *Suppose that $C_p(X)$ is strongly M -dominated for some separable metric space M . Must X be countable?*
- 3.11. Problem.** *Suppose that X is compact and $C_p(X)$ is strongly dominated by a second countable space. Is it true in ZFC that X must be countable?*
- 3.12. Problem.** *Suppose that X is a compact space and $X^2 \setminus \Delta$ is \mathbb{P} -dominated. Is it true in ZFC that X must have a small diagonal?*
- 3.13. Problem.** *Suppose that a separable metrizable space X is \mathbb{Q} -dominated. Must X be analytic?*
- 3.14. Problem.** *Suppose that every subspace of a space X is dominated by a second countable space. Is it true in ZFC that X must be cosmic?*
- 3.15. Problem.** *Suppose that every subspace of a space X is \mathbb{Q} -dominated. Is it true in ZFC that X must be cosmic?*
- 3.16. Problem.** *Suppose that every subspace of a space X is strongly dominated by a second countable space. Is it true in ZFC that X must be an \aleph_0 -space?*
- 3.17. Problem.** *Suppose that every subspace of a compact space X is dominated by a second countable space. Is it true in ZFC that X must be metrizable?*

3.18. Problem. Suppose that X is a compact space and every subspace of X is \mathbb{Q} -dominated. It is true in ZFC that X must be metrizable?

3.19. Problem. Suppose that every subspace of a compact space X is strongly dominated by a second countable space. Is it true in ZFC that X must be metrizable?

3.20. Problem. Suppose that X is a compact space and every subspace of X is strongly \mathbb{Q} -dominated. It is true in ZFC that X must be metrizable?

REFERENCES

- [Ar1] A.V. Arhangel'skii, *Structure and classification of topological spaces and cardinal invariants (in Russian)*, Uspehi Mat. Nauk, **33:6**(1978), 29-84.
- [Ar2] A.V. Arhangel'skii, *Topological Function Spaces*, Kluwer Acad. Publ., Dordrecht, 1992.
- [Ar3] A.V. Arhangel'skii, *C_p -theory*, Modern Progress in General Topology, ed. by M. Hušek and J. van Mill, Elsevier Science Publishers, B.V., Amsterdam, 1992.
- [Bat] D.P. Baturov, *On subspaces of function spaces (in Russian)*, Vestnik MGU, Matematika, Mech., **42:4**(1987), 66–69.
- [Ca] B. Cascales, *On K -analytic locally convex spaces*, Arch., Math., **49**(1987), 232–244.
- [CKS] B. Cascales, J. Kąkol, S.A. Saxon, *Weight of precompact subsets and tightness*, J., Math. Anal. Appl., **269**(2002), 500–518.
- [CMO] B. Cascales M. Muñoz and J. Orihuela, *Number of K -determination of topological spaces*, preprint.
- [CO1] B. Cascales and J. Orihuela, *On compactness in locally convex spaces*, Math. Z., **195**(1987), 365–381.
- [CO2] B. Cascales and J. Orihuela, *A sequential property of set-valued maps*, J. Math. Analysis. and Appl., **156:1**(1991), 86–100.
- [Chr] J.P.R. Christensen, *Topology and Borel Structure*, North Holland, Amsterdam, Math. Studies **10**(1974).
- [En] R. Engelking, *General Topology*, PWN, Warszawa 1977.
- [FS1] J.C. Ferrando, J. Kąkol, M. López Pellicer and S.A. Saxon, *Tightness and distinguished Fréchet spaces*, J. Math. Anal. Appl., **324:2**(2006), 862–881.
- [FS2] J.C. Ferrando, J. Kąkol, M. López Pellicer and S.A. Saxon, *Quasi-Suslin weak duals*, J. Math. Anal. Appl., **339:2**(2008), 1253–1263.
- [Ho] R.E. Hodel, *On a theorem of Arhangel'skii concerning Lindelöf p -spaces*, Canadian J. Math., **27:2**(1975), 459–468.
- [Ju] I. Juhász, *Cardinal Functions in Topology — Ten Years Later*, Mathematical Centre Tracts, **123**, Amsterdam, 1980.
- [JuS] I. Juhász, Z. Szentmiklóssy, *Convergent free sequences in compact spaces*, Proceedings of the Amer. Math. Soc., **116:4**(1992), 1153–1160.
- [KS] J. Kąkol, S.A. Saxon, *Montel (DF)-spaces, sequential (LM)-spaces and the strongest locally convex topology*, J. London Math. Soc. (2), **66:2**(2002), 388–406.

- [Mi] E. Michael, \aleph_0 -spaces, Journal of Math. and Mech., **15:6**(1966), 983–1002.
- [MS] S. Mazurkiewicz, W. Sierpiński, *Contribution à la topologie des ensembles dénombrables*, Fund. Math., **1**(1920), 17–27.
- [Mu] M. Muñoz, *Índice de K-determinación de Espacios Topológicos y σ -Fragmentabilidad de Aplicaciones*, PhD Thesis, Department of Mathematics, Murcia University, Murcia, Spain.
- [Ok] O.G. Okunev, *On Lindelöf Σ -spaces of continuous functions in the pointwise topology*, Topology Appl., **49:2**(1993), 149-166.
- [Or] J. Orihuela, *Pointwise compactness in spaces of continuous functions*, J. London Math. Soc. (2), **36:1**(1987), 143–152.
- [RJ] C.A. Rogers and J.E. Jayne, *K-analytic sets*, Analytic Sets, Academic Press Inc., London, 1980, 1-181.
- [Ro] J. Roitman, *Basic S and L*, Handbook of Set-Theoretic Topology, ed. by K. Kunen and J.E. Vaughan, North Holland, Amsterdam, 1984, 295–326.
- [Ta] M. Talagrand, *Espaces de Banach faiblement K-analytiques*, Ann. of Math., **110**(1979), 407–438.
- [Tk] V.V. Tkachuk, *A space $C_p(X)$ is dominated by irrationals if and only if it is K-analytic*, Acta Math. Hung., **107(4)**(2005), 253-265.

CASCALES SALINAS, BERNARDO

Departamento de Matemáticas,
 Facultad de Ciencias,
 Universidad de Murcia, 30.100
 Espinardo, Murcia, Spain
 e-mail: beca@um.es

ORIHUELA CALATAYUD, JOSÉ

Departamento de Matemáticas,
 Facultad de Ciencias,
 Universidad de Murcia, 30.100
 Espinardo, Murcia, Spain
 e-mail: joseori@um.es

TKACHUK, VLADIMIR V.

Departamento de Matemáticas,
 Universidad Autónoma Metropolitana,
 Av. San Rafael Atlixco, 186, Col. Vicentina
 Iztapalapa, C.P. 09340, México D.F.
 e-mail: vova@xanum.uam.mx